



NEW YORK UNIVERSITY

Institute of Mathematical Sciences

Division of Electromagnetic Research

RESEARCH REPORT No. BR-29

The Influence of Edges and Corners on Potential Functions of Surface Layers

ROLF LEIS

Mathematics Division

Air Force Office of Scientific Research

Contract No. AF 49(638)229

Project No. 47500

JUNE, 1959

104-34
21

NEW YORK UNIVERSITY
Institute of Mathematical Sciences
Division of Electromagnetic Research
Research Report No. BR-29

The Influence of Edges and Corners
on Potential Functions of Surface Layers

Rolf Leis

Rolf Leis

Rolf Leis

Sidney Borowitz
Sidney Borowitz
Acting Project Director

Qualified requestors may obtain copies of this report from the ASTIA Document Service Center, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services or have their 'need-to-know' certified by the cognizant military agency of their project or contract.

New York, June 1959

Abstract

The potential functions of multiple surface layers behave singularly when approaching the boundary of the surface. The degree of the singularities of these functions and their derivatives are discussed in this paper.

Table of Contents

	<u>Page</u>
1. Introduction	1
2. The behavior of potential functions of single and double layers	9
3. The behavior of potential functions of multiple layers	12
4. The singular behavior of potential functions of curve layers	16
5. The reducing of the differentiability suppositions	20
References	21

Let S be an analytic, orientable surface in the three dimensional space; let p and q be two points, $r(p,q) = |p-q|$ the distance between p and q , and $\rho(q)$ a function defined for all $q \in S$. With $\frac{\partial}{\partial n_q}$ we denote the derivative in the normal direction relative to $q \in S$. The potential function of the N -fold layer $\rho(q)$ is then given by

$$U_N(p) = \int_S \rho(q) \left(\frac{\partial}{\partial n_q} \right)^N \frac{1}{r(p,q)} dS_q .$$

The functions $U_N(p)$ are analytic in the neighborhood of all points p , not lying on S . $U_N(p)$ jumps, passing with p , through an interior point of S . Papers by Liapounoff, Poincaré' and E. Schmidt (cf. [1]) deal with these jump relations. They are explicitly stated for potential functions and their derivatives of arbitrary order in a paper by C. Müller^[2]. However, these papers deal only with the behavior of potential functions approaching, with p , an interior point of the surface. But for many problems it is of interest to know the behavior of potential functions and their derivatives when approaching, with p , the boundary C of the surface S . This behavior will be investigated here. We shall see that the potential functions behave singularly when approaching, with p , the boundary C . The degree of the singularity will be given explicitly.

In order not to initially burden the representation with differentiability suppositions, let us for the present assume the surface S and the layer function S to be analytic and the boundary curve C to be piecewise analytic. This assumption will be restricted to differentiability suppositions of finite order in the last section.

The differential geometric formalism which we need will be given in the first section where we follow the notations given in C. Müller's paper [2]. The second section deals with the functions U_0 , U_1 and their gradients. In the third section, the singular behavior of the functions U_N and ∇U_N will be reduced to curve integrals over the boundary C . These curve integrals will be discussed in Section 4. Section 5 finally restricts the differentiability suppositions.

1. Introduction

Let the orientable surface S be bounded by the piecewise analytic Jordan curve C . Let S have the following properties:

1. There exists a coordinate system for every point P of S (a closed domain).
2. P is the origin of the coordinate system.
3. A constant $c > 0$ exists such that the part of S lying in the interior of the sphere $(x^1)^2 + (x^2)^2 + (x^3)^2 \leq c$ can be represented in the form $x^3 = \phi(x^1, x^2)$. $\phi(x^1, x^2)$ is an analytic function for all $(x^1)^2 + (x^2)^2 < c$.
4. The derivatives $\frac{\partial \phi}{\partial x^1} = \phi_{|1}$, $\phi_{|2}$ and $\phi_{|1|2}$ vanish in the origin.

Thus the x^3 -axis is normal to S in P ; the x^1 - and x^2 -axis are directions of principal curvature. Therefore, we briefly call this coordinate system a tangent-normal system.

Let e_i be the unit vectors in the direction of the coordinate axis;

$$(1.1) \quad x^1 e_1 + x^2 e_2 + \phi(x^1, x^2) e_3 = f(x^1, x^2)$$

then represents a point of the surface. f denotes a vector with the components f^1, f^2, f^3 .

$$(1.2) \quad f_{|\mu} \stackrel{\circ}{=} e_{\mu} ; \quad f_{|1|2} = \phi_{|1|2} e_3 \stackrel{\circ}{=} 0.$$

In (1.2) and in the following equations, $\stackrel{\circ}{=}$ denotes equal in the origin (in P); Greek subscripts stand for 1,2; Roman for 1,2,3.

Now we introduce a new coordinate system - in short a u-system - by

$$(1.3) \quad x = f(u^1, u^2) + u^3 n(u^1, u^2)$$

where n denotes the unit vector orthogonal to the surface. Then we have

Then we have

$$(1.4) \quad \frac{\partial x^i}{\partial u^x} \stackrel{\circ}{=} \delta_x^i$$

and with

$$(1.5) \quad n_{|\mu} = -L_{\mu}^{\sigma} f_{|\sigma} ; \quad \gamma_{\mu\nu} = (f_{|\mu} - u^3 L_{\mu}^{\sigma} f_{|\sigma}) (f_{|\nu} - u^3 L_{\nu}^{\sigma} f_{|\sigma}) ,$$

$$(1.6) \quad \left\{ \begin{aligned} g_{\mu\nu} &= (x_{|\mu} x_{|\nu}) = (f_{|\mu} - u^3 L_{\mu}^{\sigma} f_{|\sigma}) (f_{|\nu} - u^3 L_{\nu}^{\sigma} f_{|\sigma}) \\ &= \gamma_{\mu\nu} - 2u^3 L_{\mu\nu} + (u^3)^2 L_{\mu}^{\sigma} L_{\sigma\nu} . \end{aligned} \right.$$

Let $2H = k_1 + k_2$ be the mean curvature and $K = k_1 k_2$, the Gauss curvature.

Then, with

$$(1.7) \quad L_{11} \stackrel{\circ}{=} k_1 ; L_{22} \stackrel{\circ}{=} k_2 ; L_{12} \stackrel{\circ}{=} 0 ; \gamma_{\mu\nu} = \delta_{\mu\nu} ,$$

we have

$$(1.8) \quad g_{\mu\nu} \stackrel{\circ}{=} (1 - (u^3)^2 K) \gamma_{\mu\nu} - 2u^3 (1 - u^3 H) L_{\mu\nu} .$$

Equation (1.8) represents a relation between tensors which is valid in the point P. However, the point P is in no way distinguished on the surface; rather, to every arbitrary point of the surface, there exists a tangent-normal system and, consequently, a u-system in which equation (1.8) is valid. Thus, (1.8) is valid in every point and we get

$$(1.9) \quad g_{\mu\nu} = (1 - (u^3)^2 K) \gamma_{\mu\nu} - 2u^3(1 - u^3 H) L_{\mu\nu}$$

and

$$(1.10) \quad g_{33} = (x_{|3} x_{|3}) = n^2 = 1; \quad g_{3\mu} = (x_{|3} x_{|\mu}) = (n(f_{| \nu} - u^3 L_{\mu}^{\sigma} f_{|\sigma})) = 0.$$

Let g be $\det g_{ik}$ and γ be $\det \gamma_{\mu\nu}$, then g/γ is an invariant of the surface (u^3 fixed) and we get

$$(1.11) \quad g \triangleq \gamma(1 - 2u^3 H + (u^3)^2 K)^2$$

or with

$$(1.12) \quad G = 1 - 2u^3 H + (u^3)^2 K$$

and

$$(1.13) \quad g = \gamma G^2.$$

We get the Delta operator

$$(1.14) \quad \Delta = \left(\frac{\partial}{\partial x^1} \right)^2 + \left(\frac{\partial}{\partial x^2} \right)^2 + \left(\frac{\partial}{\partial x^3} \right)^2$$

in the u-system by

$$(1.15) \quad \Delta = \frac{1}{\sqrt{g}} \frac{\partial}{\partial u^i} \sqrt{g} g^{ik} \frac{\partial}{\partial u^k}$$

or

$$(1.16) \quad \Delta = \frac{1}{G} \frac{1}{\sqrt{\gamma}} \frac{\partial}{\partial u^\mu} G \sqrt{\gamma} g^{\mu\nu} + \frac{1}{G} \frac{\partial}{\partial u^3} G \frac{\partial}{\partial u^3} .$$

For sufficiently small u^3 we expand

$$(1.17) \quad G g^{\mu\nu} = \sum_{j=0}^{\infty} \frac{(u^3)^j}{j!} S_{(j)}^{\mu\nu}$$

and denote

$$(1.18) \quad \Delta_j = \frac{1}{\sqrt{\gamma}} \frac{\partial}{\partial u^\mu} \sqrt{\gamma} S_{(j)}^{\mu\nu} \frac{\partial}{\partial u^\nu} \quad \left(\Delta_0 = \frac{1}{\sqrt{\gamma}} \frac{\partial}{\partial u^\mu} \sqrt{\gamma} g^{\mu\nu} \frac{\partial}{\partial u^\nu} \right) .$$

Thus we get

Lemma 1: For sufficiently small u^3 we have

$$G \Delta = \sum_{j=0}^{\infty} \frac{(u^3)^j}{j!} \Delta_j + \frac{\partial}{\partial u^3} G \frac{\partial}{\partial u^3} .$$

From Lemma 1, it follows for $u^3 = 0$ and for harmonic functions with

$$(1.19) \quad \frac{\partial}{\partial u^3} = \frac{\partial}{\partial n} \quad \text{on } S$$

$$(1.20) \quad \Delta_0 U + \left(\frac{\partial^2}{\partial n^2} - 2H \frac{\partial}{\partial n} \right) U = 0 .$$

In analogy to Lemma 1, we get, by differentiation with respect to u^3 :

Lemma 2: Let the function U be harmonic in the neighborhood of a point P .

Then the derivatives of U satisfy the recurrence relations

$$0 = \left\{ \sum_{j=0}^N \binom{N}{j} \Delta_{N-j} \left(\frac{\partial}{\partial n} \right)^j + \left(\frac{\partial}{\partial n} \right)^{N+2} - 2H(N+1) \left(\frac{\partial}{\partial n} \right)^{N+1} + 2K \binom{N+1}{2} \left(\frac{\partial}{\partial n} \right)^N \right\} U.$$

We get the Nabla operator

$$(1.21) \quad \nabla = \frac{\partial}{\partial x^1} e_1 + \frac{\partial}{\partial x^2} e_2 + \frac{\partial}{\partial x^3} e_3$$

in the u -system by

$$(1.22) \quad \nabla U = g^{ki} x_{|i} U_{|k} = g^{\mu\nu} (f_{|\nu} + u^3 n_{|\nu}) U_{|\mu} + n U_{|3}.$$

We develop for small values of u^3

$$(1.23) \quad g^{\mu\nu} (f_{|\nu} + u^3 n_{|\nu}) = g^{\mu\nu} (\delta_{\nu}^{\rho} - u^3 L_{\nu}^{\rho}) f_{|\rho}$$

$$= f_{|\rho} \sum_{j=0}^{\infty} \frac{(u^3)^j}{j!} Q_{(j)}^{\mu\rho}$$

and use the notation

$$(1.24) \quad \nabla_j^{\mu\nu} = Q_{(j)}^{\mu\nu} f_{|\nu} \frac{\partial}{\partial u^{\mu}} \quad \left(\nabla_o = \gamma^{\mu\nu} f_{|\nu} \frac{\partial}{\partial u^{\mu}} \right).$$

Then it follows:

Lemma 3: For sufficiently small u^3 , we have

$$\nabla = \sum_{j=0}^{\infty} \frac{(u^3)^j}{j!} \nabla_j + n \frac{\partial}{\partial u^3}$$

Finally, we want to derive some formulas from Gauss' theorem which we shall need later on. Let J be a continuously differentiable function defined on S . Then we get for a sufficiently small surface S' with the boundary C'

$$\begin{aligned}
 (1.25) \quad \int_{S'} \nabla_j J \, dS &= \int_{S'} Q_{(j)}^{\mu\nu} f_{\mu\nu} J_{,\nu} \sqrt{\gamma} \, du^1 \, du^2 \\
 &= - \int_{S'} J \frac{\partial}{\partial u^\mu} \left(Q_{(j)}^{\mu\nu} f_{\mu\nu} \sqrt{\gamma} \right) du^1 \, du^2 \\
 &\quad + \int_{C'} \sqrt{\gamma} J f_{\mu\nu} \left(Q_{(j)}^{1\nu} \dot{u}^2 - Q_{(j)}^{2\nu} \dot{u}^1 \right) dc.
 \end{aligned}$$

Since

$$(1.26) \quad n_j = f_{\mu\nu} \left(Q_{(j)}^{1\nu} \dot{u}^2 - Q_{(j)}^{2\nu} \dot{u}^1 \right) \sqrt{\gamma} \quad ; \quad p_j = \frac{1}{\sqrt{\gamma}} \frac{\partial}{\partial u^\mu} \left(Q_{(j)}^{\mu\nu} f_{\mu\nu} \sqrt{\gamma} \right),$$

we find, after summation over S' ,

Lemma 4: Let J be a continuously differentiable function defined on S .

Then we have

$$\int_S \nabla_j J \, dS = - \int_S J p_j \, dS + \int_C J n_j \, dc.$$

For $j = 0$ we have from Lemma 4.

Lemma 4*: Let J be a continuously differentiable function defined in S .

Then we have

$$\int_S \nabla_o J \, dS = -2 \int_S n_{HJ} \, dS + \int_C n_o J \, dc$$

where n_o is a vector normal to C and orthogonal to n .

Let \mathcal{X} and \mathcal{L} be two times continuously differentiable functions defined on S . Then we have

$$\begin{aligned} (1.27) \quad \int_{S'} \mathcal{X} \Delta_j \mathcal{L} \, dS &= \int_{S'} \mathcal{X} \frac{\partial}{\partial u^\mu} (\sqrt{\gamma} s_{(j)}^{\mu\nu} \mathcal{L}_{\nu}{}^{1\nu}) du^1 du^2 \\ &= - \int_{S'} \mathcal{X}_{;\mu} \sqrt{\gamma} s_{(j)}^{\mu\nu} \mathcal{L}_{\nu}{}^{1\nu} du^1 du^2 + \int_{C'} \sqrt{\gamma} \mathcal{X} \mathcal{L}_{\nu}{}^{1\nu} (s_{(j)}^{1\nu} \dot{u}^2 - s_{(j)}^{2\nu} \dot{u}^1) \, dc \\ &= \int_{S'} \mathcal{L} \Delta_j \mathcal{X} \, dS + \int_{C'} \mathcal{X} \sqrt{\gamma} \mathcal{L}_{\nu}{}^{1\nu} (s_{(j)}^{1\nu} \dot{u}^2 - s_{(j)}^{2\nu} \dot{u}^1) \, dc \\ &\quad - \int_{C'} \mathcal{L} \sqrt{\gamma} \mathcal{X}_{;\mu} (s_{(j)}^{\mu 1} \dot{u}^2 - s_{(j)}^{\mu 2} \dot{u}^1) \, dc . \end{aligned}$$

Since

$$(1.28) \quad s_{(j)}^\nu = \sqrt{\gamma} (s_{(j)}^{1\nu} \dot{u}^2 - s_{(j)}^{2\nu} \dot{u}^1) ; s_j = s_{(j)}^{\nu} f_{\nu}{}^{1\nu} ; \nabla_o \mathcal{L} = \mathcal{L}_{\nu}{}^{1\nu} f_{\mu}{}^{1\nu} \gamma^{\mu\nu}$$

we find

Lemma 5: Let the functions \mathcal{K} and \mathcal{L} be two times continuously differentiable on S . Then we have

$$\int_S \mathcal{K} \Delta_j \mathcal{L} \, dS = \int_S \mathcal{L} \Delta_j \mathcal{K} \, dS + \int_C (\mathcal{K} \nabla_{\circ} \mathcal{L} - \mathcal{L} \nabla_{\circ} \mathcal{K}) s_j \, dc$$

or, in particular,

Lemma 5*: Let the functions \mathcal{K} and \mathcal{L} be two times continuously differentiable on S . Then we have

$$\int_S \mathcal{K} \Delta_{\circ} \mathcal{L} \, dS = \int_S \mathcal{L} \Delta_{\circ} \mathcal{K} \, dS + \int_C (\mathcal{K} n_{\circ} \nabla_{\circ} \mathcal{L} - \mathcal{L} n_{\circ} \nabla_{\circ} \mathcal{K}) \, dc$$

2. The behavior of potential functions arising from single and double layers

The potential function of the single layer $\rho(q)$

$$(2.1) \quad U_{\circ}(p) = \int_S \rho(q) \frac{1}{r(p, q)} \, dS_q$$

is continuous everywhere^[1]. The behavior of the potential function of the double layer $\mu(q)$

$$(2.2) \quad U_1(p) = \int_S \mu(q) \frac{\partial}{\partial n_q} \frac{1}{r(p, q)} \, dS_q$$

approaching, with p , an interior point of the surface S , is characterized

by the jump relations [1], [2]

$$(2.3) \quad \begin{cases} U_1|_{\text{ext.}} = 2\pi\mu + \int_S \mu \frac{\partial}{\partial n} \frac{1}{r} dS \Big|_{p \in S} \\ U_1|_{\text{int.}} = -2\pi\mu + \int_S \mu \frac{\partial}{\partial n} \frac{1}{r} dS \Big|_{p \in S} \end{cases} .$$

The integral

$$(2.4) \quad \int_S \mu(q) \frac{\partial}{\partial n} \frac{1}{r(p, q)} dS_q \Big|_{p \in S}$$

represents a function continuous for all $p \in S$. Thus $U_1(p)$ is not continuous, passing with p through S , but it is bounded for all p .

The gradients of these functions become singular, however, when approaching, with p , a point of the boundary C . To show this, we form, according to Lemma 3,

$$(2.5) \quad \nabla U_0 = - \int_S \rho(q) \nabla_q \frac{1}{r(p, q)} dS_q = - \int_S \rho n \frac{\partial}{\partial n} \frac{1}{r} dS - \int_S \rho \nabla_0 \frac{1}{r} dS.$$

From this, according to Lemma 4*, we get

$$(2.6) \quad \nabla U_0 = - \int_S \rho n \frac{\partial}{\partial n} \frac{1}{r} dS + \int_S (\nabla_0 \rho) \frac{1}{r} dS + 2 \int_S n H \rho \frac{1}{r} dS - \int_C n_0 \rho \frac{1}{r} dc .$$

The curve integral in (2.6) becomes logarithmically singular [3], whereas the other surface integrals behave regularly. Thus we get

Theorem 1: The potential function $U_0(p)$ behaves regularly approaching, with p , the boundary C , whereas $\nabla U_0(p)$ becomes logarithmically singular.

We shall now discuss $\nabla U_1(p)$ and, in a similar way, get

$$(2.7) \quad -\nabla U_1 = \int_S \mu \frac{\partial}{\partial n} \nabla_q \frac{1}{r(p, q)} dS_q = \int_S \mu n \frac{\partial^2}{\partial n^2} \frac{1}{r} dS \\ + \int_S \mu \nabla_o \frac{\partial}{\partial n} \frac{1}{r} dS + \int_S \mu \nabla_1 \frac{1}{r} dS \quad .$$

We discuss the resulting surface integrals individually. According to (1.20) and Lemma 5*, we get

$$(2.8) \quad \int_S \mu n \frac{\partial^2}{\partial n^2} \frac{1}{r} dS = 2 \int_S \mu H n \frac{\partial}{\partial n} \frac{1}{r} dS - \int_S \mu \Delta_o \frac{1}{r} dS \\ = 2 \int_S \mu H n \frac{\partial}{\partial n} \frac{1}{r} dS - \int_S (\Delta_o \mu n) \frac{1}{r} dS \\ - \int_C \left\{ \mu n (n_o \nabla_o) \frac{1}{r} - \frac{1}{r} (n_o \nabla_o) \mu n \right\} dc \quad .$$

The curve integral behaves singularly, like $\frac{1}{r}$, as p approaches $C[3]$.

According to Lemma 4*, we get, for the second integral in (2.7),

$$(2.9) \quad \int_S \mu \nabla_o \frac{\partial}{\partial n} \frac{1}{r} dS = - \int_S (\nabla_o \mu) \frac{\partial}{\partial n} \frac{1}{r} dS - 2 \int_S n H \mu \frac{\partial}{\partial n} \frac{1}{r} dS \\ + \int_C n_o \mu \frac{\partial}{\partial n} \frac{1}{r} dc \quad .$$

The curve integral in (2.9) also behaves singularly, like $\frac{1}{r}$. Finally, for the third integral in (2.7), according to Lemma 4, we have

$$(2.10) \quad \int_S \mu \nabla_1 \frac{1}{r} dS = - \int_S (\nabla_1 \mu) \frac{1}{r} dS - \int_S \mu \frac{1}{r} p_1 dS + \int_C \mu \frac{1}{r} n_0 dc .$$

Again the curve integral becomes logarithmically singular. Thus we get

Theorem 2: The potential function U_1 is bounded, approaching, with p , the boundary C , whereas ∇U_1 becomes singular like $\frac{1}{r}$ (r being the shortest distance between p and C).

3. The behavior of potential functions of multiple layers

In the last section, we discussed the behavior of potential functions of single and double layers and their derivatives. Now we want to extend these discussions to the potential functions of N -fold layers

$$(3.1) \quad U_N(p) = \int_S \rho(q) \left(\frac{\partial}{\partial n_q} \right)^N \frac{1}{r(p, q)} dS_q .$$

To do this, we first of all want to express the operator $\left(\frac{\partial}{\partial n} \right)^N$ through lower derivatives. We make the assumption

$$(3.2) \quad \left(\frac{\partial}{\partial n} \right)^N = \Omega_N^1 + \Omega_N^2 \frac{\partial}{\partial n} ,$$

and, according to Lemma 2, get the recursions

$$(3.3) \quad \Omega_{N+2}^\mu = 2H(N+1) \Omega_{N+1}^\mu - 2K \binom{N+1}{2} \Omega_N^\mu - \sum_{j=0}^N \binom{N}{j} \Delta_{N-j} \Omega_j^\mu$$

with the initial values

$$(3.4) \quad \Omega_0^1 = 1 \quad ; \quad \Omega_0^2 = 0 \quad ; \quad \Omega_1^1 = 0 \quad ; \quad \Omega_1^2 = 1 \quad .$$

It follows from (3.3) and (3.4) that the operators Ω_{2N}^1 and Ω_{2N+1}^1 contain derivatives of maximal order $2N$ and that the operators Ω_{2N-1}^2 and Ω_{2N}^2 contain derivatives of maximal order $2N-2$.

Now we form the adjoint operators Φ_N^μ to Ω_N^μ according to

$$(3.5) \quad \int_S \mathcal{X} \Omega_N^\mu \mathcal{L} \, dS = \int_S \mathcal{L} \Phi_N^\mu \mathcal{X} \, dS + \int_C \chi_N^\mu (\mathcal{X}, \mathcal{L}) \, dc$$

where \mathcal{X} and \mathcal{L} denote sufficiently differentiable functions. As the operators Δ_{N-j} are self-adjoint, we get, for Φ_N^μ , the recursion

$$(3.6) \quad \Phi_{N+2}^\mu = 2(N+1) \Phi_{N+1}^\mu H - 2 \binom{N+1}{2} \Phi_N^\mu K - \sum_{j=0}^N \binom{N}{j} \Phi_j^\mu \Delta_{N-j}$$

with the initial values

$$(3.7) \quad \Phi_0^1 = 1 \quad ; \quad \Phi_0^2 = 0 \quad ; \quad \Phi_1^1 = 0 \quad ; \quad \Phi_1^2 = 1 \quad .$$

According to Lemma 5, we get, for the χ_N^μ ,

$$(3.8) \quad \begin{aligned} \chi_{N+2}^\mu &= 2(N+1) \chi_{N+1}^\mu (H \mathcal{X}, \mathcal{L}) - 2 \binom{N+1}{2} \chi_N^\mu (K \mathcal{X}, \mathcal{L}) \\ &- \sum_{j=0}^N \binom{N}{j} \left\{ (\mathcal{X} \nabla_0 \Omega_j^\mu \mathcal{L} - (\Omega_j^\mu \mathcal{L}) \nabla_0 \mathcal{X}) s_{n-j} \right. \\ &\left. + \chi_j^\mu (\Delta_{N-j} \mathcal{X}, \mathcal{L}) \right\} \end{aligned}$$

with the initial values.

$$(3.9) \quad x_0^1 = x_1^1 = x_0^2 = x_1^2 = 0.$$

From (3.8) and (3.9) it follows that $x_{2N}^1(\mathcal{X}, \mathcal{L})$ and $x_{2N+1}^1(\mathcal{X}, \mathcal{L})$ contain derivatives of maximal order $2N-1$ with reference to \mathcal{L} , and that $x_{2N-1}^2(\mathcal{X}, \mathcal{L})$ and $x_{2N}^2(\mathcal{X}, \mathcal{L})$ contain derivatives of maximal order $2N-3$ with reference to \mathcal{L} . $x_{2N}^\mu(\mathcal{X}, \mathcal{L})$ and $x_{2N+1}^\mu(\mathcal{X}, \mathcal{L})$ contain derivatives of maximal order $2N-1$ with reference to \mathcal{X} .

Thus we get

Lemma 6: Let the function \mathcal{X} be analytic and \mathcal{L} be harmonic. Then there exist operators Ω_N^μ , Φ_N^μ , χ_N^μ defined in (3.3, 6, 8) with

$$\left(\frac{\partial}{\partial n}\right)^N \mathcal{L} = \left(\Omega_N^1 + \Omega_N^2 \frac{\partial}{\partial n}\right) \mathcal{L}$$

and

$$\int_S \mathcal{X} \Omega_N^\mu \mathcal{L} \, dS = \int_S \mathcal{L} \Phi_N^\mu \mathcal{L} \, dS + \int_C \chi_N^\mu(\mathcal{L}, \mathcal{X}) \, dc.$$

It follows from Lemma 6 that

$$\begin{aligned} (3.10) \quad U_N &= \int_S \rho \left(\frac{\partial}{\partial n}\right)^N \frac{1}{r} \, dS = \int_S \rho \Omega_N^1 \frac{1}{r} \, dS + \int_S \rho \Omega_N^2 \frac{\partial}{\partial n} \frac{1}{r} \, dS \\ &= \int_S \left(\Phi_N^1 \rho\right) \frac{1}{r} \, dS + \int_S \left(\Phi_N^2 \rho\right) \frac{\partial}{\partial n} \frac{1}{r} \, dS \\ &\quad + \int_C \chi_N^1\left(\rho, \frac{1}{r}\right) \, dc + \int_C \chi_N^2\left(\rho, \frac{\partial}{\partial n} \frac{1}{r}\right) \, dc. \end{aligned}$$

Thus, we have expressed the function U_N through functions of the type U_0 and U_1 and boundary integrals. These curve integrals in (3.10) therefore describe the singular behavior of the functions $U_N(p)$ approaching, with p , the boundary C . We shall discuss the singular behavior of these integrals in the next section.

For the gradient ∇U_N we get, according to Lemmas 3 and 4,

$$\begin{aligned}
 (3.11) \quad -\nabla U_N &= \int_S \rho \left(\frac{\partial}{\partial n} \right)^N \nabla \frac{1}{r} \, dS = \int_S \rho n \left(\frac{\partial}{\partial n} \right)^{N+1} \frac{1}{r} \, dS \\
 &+ \sum_{j=0}^N \int_S \rho \nabla_{N-j} \left(\frac{\partial}{\partial n} \right)^j \, dS \\
 &= \int_S \rho n \left(\frac{\partial}{\partial n} \right)^{N+1} \frac{1}{r} \, dS - \sum_{j=0}^N \int_S \left(\nabla_{N-j} \rho \right) \left(\frac{\partial}{\partial n} \right)^j \frac{1}{r} \, dS \\
 &- \sum_{j=0}^N \left\{ \int_S p_j \rho \left(\frac{\partial}{\partial n} \right)^j \frac{1}{r} \, dS - \int_C n_j \rho \left(\frac{\partial}{\partial n} \right)^j \frac{1}{r} \, dc \right\}.
 \end{aligned}$$

The surface integrals in (3.11) are again functions of the form U_K . We have now expressed their singular behavior in terms of curve integrals which we shall discuss in the next paragraph.

4. The singular behavior of potential functions arising from one-dimensional layers

In the last section we expressed the singular behavior of potential functions of surface layers through curve integrals over the boundary C of the surface S . We shall now discuss these integrals. As the derivatives with respect to all coordinates u^i enter into the integrands, we shall discuss in particular

$$(4.1) \quad V_N = \int_C \rho \left(\frac{\partial}{\partial c} \right)^N \frac{1}{r} dc$$

and

$$(4.2) \quad W_N = \int_C \rho \left(\frac{\partial}{\partial n} \right)^N \frac{1}{r} dc$$

where c denotes the arclength and n a vector normal to C .

We assumed C to be piecewise analytic, so let us assume that $C = \sum_{i=1}^K C_i$, where the C_i are analytic curves. Through partial integration, it follows from (4.1) that

$$(4.3) \quad V_N = \sum_i \int_{C_i} \rho \left(\frac{\partial}{\partial c_i} \right)^N \frac{1}{r} dc = \sum_i \left\{ \left[\rho \left(\frac{\partial}{\partial c_i} \right)^{N-1} \frac{1}{r} \right]_0^{L_i} - \int_{C_i} \left(\frac{\partial}{\partial c_i} \right) \rho \left(\frac{\partial}{\partial c_i} \right)^{N-1} \frac{1}{r} dc_i \right\}$$

or

$$(4.4) \quad V_N = \sum_i \left\{ - \sum_{j=1}^N (-1)^j \left[\left(\frac{\partial}{\partial c_i} \right)^{j-1} \rho \left(\frac{\partial}{\partial c_i} \right)^{N-j} \frac{1}{r} \right]_0^{L_i} + (-1)^N \int_{C_i} \left(\frac{\partial}{\partial c_i} \right)^N \rho \frac{1}{r} dc_i \right\}.$$

Thus we get [3]

Lemma 7: The function $V_N(p)$ becomes logarithmically singular when approaching C , with p . For $N \geq 1$ there are additional singularities at the corners of C of the order r^{-N} (r being the distance from p to a corner).

$$\begin{aligned}
 (4.6) \quad |W_1^{\tau_1}| &\leq A \left| \int_{-\tau_1}^{\tau_1} \frac{\bar{n} \bar{r}}{r^3} dx \right| = O \left(\int_{-\tau_1 - c_1}^{\tau_1 - c_1} \frac{x^2 + x(y - y_1) + (z - z_1)}{\sqrt{x^2 + (y - y_1)^2 + (z - z_1)^2}^3} dx \right) \\
 &= O \left(\int_{-\tau_1 - c_1}^{\tau_1 - c_1} \frac{dx}{r^2} \right).
 \end{aligned}$$

With

$$(4.7) \quad r = \sqrt{x^2 + (y - y_1)^2 + (z - z_1)^2} \quad ; \quad \rho = \sqrt{y_1^2 + z_1^2} \quad ; \quad R = \sqrt{x^2 + \rho^2}$$

we get

$$\begin{aligned}
 (4.8) \quad \left| \frac{1}{r} - \frac{1}{R} \right| &= \frac{|R^2 - r^2|}{rR(r + R)} = \frac{|y|(|y - y_1| + |y_1|) + |z|(|z - z_1| + |z_1|)}{rR(r + R)} \\
 &= O(1)
 \end{aligned}$$

and, by induction, it follows that

$$(4.9) \quad \left| \frac{1}{r^N} - \frac{1}{R^N} \right| = O\left(\frac{1}{R^{N-1}}\right).$$

Thus we get

$$(4.10) \quad |W_1^{\tau_1}| = O\left(\int \frac{dx}{R^2}\right) = O\left(\frac{1}{\rho}\right),$$

ρ being the distance of Q from the curve C (in the normal plane). The example of the straight line shows that in general the estimate in (4.10) cannot be improved.

In a similar way, we get, in the general case ($N \geq 1$)

$$(4.11) \quad |W_N^1| = O\left(\int \frac{dx}{R^{N+1}}\right) = O\left(\frac{1}{\rho^N}\right) .$$

Thus we have

Lemma 8: The function $W_N(\rho)$ becomes singular, like ρ^{-N} , if p approaches the curve C . ρ is the distance from p to the curve C taken in the normal plane of the curve.

We can now describe the singular behavior of the functions

$$(4.12) \quad U_N = \int_S \rho \left(\frac{\partial}{\partial n}\right)^N \frac{1}{r} dS .$$

As the curve integrals in (3.10) contain derivatives of $\frac{1}{r}$ of the maximal order $N-1$, we get

Theorem 3: The potential functions

$$U_N(p) = \int_S \rho \left(\frac{\partial}{\partial n}\right)^N \frac{1}{r} dS \quad N > 1$$

become singular, like ρ^{-N} , if p approaches the curve C (ρ being the distance from p to the curve C taken in the normal plane of the curve): Besides this, there are point singularities at the corners of order ρ_1^{1-N} (ρ_1 being the distance from p to the corner).

Similarly, from (3.11), we get

Theorem 4: The potential functions

$$\nabla U_N = \nabla \int_S \rho \left(\frac{\partial}{\partial n} \right)^N \frac{1}{r} dS \quad N \geq 1$$

become singular, like ρ^{-N} , if p approaches the curve C . There are point singularities at the corners of order ρ_1^{-N} .

5. The reduction of the differentiability assumptions

In order to simplify the presentation, we have so far assumed an analytic behavior for the surface S , for the boundary curve C and for the layer function ρ . However, this analytic behavior was not used. Actually, it is sufficient to make differentiability assumptions of finite order. While discussing the behavior of the functions U_N in interior points of S these assumptions are stated in [2]. Thus, let us confine ourselves to boundary points. In discussing the functions U_N , only derivative of order $N-1$ occurred. Thus, we assume C, S and ρ to be $(N-1)$ -times differentiable. To be able to introduce the tangent-normal coordinate system, however, C has to be at least twice (or one-times Hölder-continuously) differentiable. The layer function ρ has at least to be bounded. For the discussion of ∇U , the derivatives of order N should exist.

References

- [1] Lichtenstein. L - Neuere Entwicklung der Potentialtheorie
 (Enzk.d.Math.Wiss.;II/3,1 p.197,ff).

- [2] Müller, C - Die Potentiale einfacher und mehrfacher
 Flächenbelegungen. Math,Ann.123 p.235 ff 1951.

- [3] Müller C. and - Über Potentialfunktionen von Kurvenbele-
 Leis R. gungen. Arch.Rat.Mech.An.2,p.87 ff 1958.

Distribution List for Research Reports

Contract No. AF 49(63R)229

(ONE copy unless otherwise noted)

(3) Commander European Office, ARDC 47 Rue Camillesteven Brussels, Belgium	Commander Air Force Missile Test Center ATTN: Technical Library Patrick Air Force Base, Florida	Department of Mathematics Washington University St. Louis 5, Missouri
Applied Mathematics and Statistics Lab. Stanford University Stanford, California	Department of Mathematics Northwestern University Evanston, Illinois	Department of Mathematics University of Missouri Columbia, Missouri
Department of Mathematics University of California Berkeley, California	Institute for Air Weapons Research Museum of Science and Industry University of Chicago Chicago 37, Illinois	Linda Hall Library ATTN: Mr. Thomas Gillis Document Division 5109 Cherry Street Kansas City 10, Missouri
Commander Air Force Flight Test Center ATTN: Technical Library Edwards Air Force Base, California	Department of Mathematics University of Chicago Chicago 37, Illinois	Commander Strategic Air Command ATTN: Operations Analysis Offutt Air Force Base Omaha, Nebraska
The Rand Corporation Technical Library 1700 Main Street Santa Monica, California	Department of Mathematics University of Illinois Urbana, Illinois	The James Forrestal Res. Center Lib. Princeton University Princeton, New Jersey
Director of Advanced Studies Air Force Office of Scientific Research Post Office Box 2035 Pasadena 2, California	Department of Mathematics Purdue University Lafayette, Indiana	Library Institute for Advanced Study Princeton, New Jersey
Commander Western Development Division ATTN: WDSIT P.O. Box 262 Inglewood, California	Institute for Fluid Dynamics and Applied Mathematics University of Maryland College Park, Maryland	Department of Mathematics Fine Hall Princeton University Princeton, New Jersey
Department of Mathematics Yale University New Haven, Connecticut	Mathematics and Physics Library The Johns Hopkins University Baltimore, Maryland	Commander AF Missile Development Center ATTN: Technical Library Holloman Air Force Base, New Mexico
Chief of Naval Operations Department of the Navy ATTN: OP03EG; Mr. Arthur Kaufman Washington 25, D.C.	Commander HQ Air Research and Development Command Director of Research Andrews Air Force Base Washington 25, D.C.	Commander Air Force Special Weapons Center ATTN: Technical Library Kirtland Air Force Base Albuquerque, New Mexico
(2) Office of Naval Research Department of the Navy ATTN: Code 432 Washington 25, D.C.	Department of Mathematics Harvard University Cambridge 38, Massachusetts	Prof. J. Wolfowitz Mathematics Department White Hall Cornell University Ithaca, New York
Department of Commerce Office of Technical Services Washington 25, D.C.	Department of Mathematics Massachusetts Institute of Technology Cambridge, Massachusetts	Department of Mathematics Syracuse University Syracuse, New York
Director National Security Agency ATTN: Dr. H.H. Campaigne Washington 25, D.C.	Commander Air Force Cambridge Research Center ATTN: Geophysics Research Library L.G. Hanscom Field Bedford, Massachusetts	New York University Institute of Mathematical Sciences Division of Electromagnetic Research ATTN: Professor M. Kline 25 Waverly Place New York 3, N.Y.
Library National Bureau of Standards Washington 25, D.C.	Commander Air Force Cambridge Research Center ATTN: Electronics Research Library L.G. Hanscom Field Bedford, Massachusetts	Department of Mathematics Columbia University ATTN: Professor B.C. Koopman New York 27, N.Y.
National Applied Mathematics Labs. National Bureau of Standards Washington 25, D.C.	Psychological Labs. ATTN: Dr. George A. Miller Memorial Hall Harvard University Cambridge 38, Mass.	Department of Mathematical Statistics Fayerweather Hall ATTN: Dr. Herbert Robbins Columbia University New York 27, N.Y.
Headquarters, USAF Assistant for Operations Analysis Deputy Chief of Staff, Operations, AFOSM Washington 25, D.C.	Department of Mathematics Wayne University ATTN: Dr. Y.W. Chen Detroit 1, Michigan	Commander Rome Air Development Center ATTN: Technical Library Griffiss Air Force Base Rome, New York
(2) Commander Air Force Office of Scientific Research ATTN: SRDB Washington 25, D.C.	Willow Run Research Center University of Michigan Ypsilanti, Michigan	Institute for Aeronautical Sciences ATTN: Librarian 2 East 60th Street New York 21, N.Y.
(2) Commander Air Force Office of Scientific Research ATTN: SRE Washington 25, D.C.	Department of Mathematics Folwell Hall University of Minnesota Minneapolis, Minnesota	
National Science Foundation ATTN: Dr. Leo Cohen Washington 25, D.C.	Department of Mathematics Institute of Technology Engineering Building University of Minnesota Minneapolis, Minnesota	
Commander Air Force Armament Center ATTN: Technical Library Eglin Air Force Base, Florida		

- Institute of Statistics
North Carolina State College of A and E
Raleigh, North Carolina
- Department of Mathematics
University of North Carolina
Chapel Hill, North Carolina
- (2) Office of Ordnance Research
Box CM
Duke Station
Durham, North Carolina
- Department of Mathematics
Duke University
Duke Station
Durham, North Carolina
- Commander
Air Technical Intelligence Center
ATTN: ATIAE-4
Wright-Patterson Air Force Base, Ohio
- Commander
Wright Air Development Center
ATTN: Technical Library
Wright-Patterson Air Force Base, Ohio
- (2) Commander
Wright Air Development Center
ATTN: ARL Technical Library, WCRR
Wright-Patterson Air Force Base, Ohio
- (2) Commandant
USAF Institute of Technology
ATTN: Technical Library, MCLI
Wright-Patterson Air Force Base, Ohio
- Department of Mathematics
University of Pennsylvania
Philadelphia, Pennsylvania
- Commander
Arnold Engineering Development Center
ATTN: Technical Library
Tullahoma, Tennessee
- Defense Research Laboratory
University of Texas
Austin, Texas
- Department of Mathematics
Rice Institute
Houston, Texas
- Commander
AF Personnel and Training Research Center
ATTN: Technical Library
Lackland Air Force Base
San Antonio, Texas
- Q10 Armed Services Technical Information Agency
Arlington Hall Station
Arlington 12, Virginia
- Department of Mathematics
University of Wisconsin
Madison, Wisconsin
- Mathematics Research Center
ATTN: R.E. Langer
University of Wisconsin
Madison, Wisconsin
- Naval Electronics Laboratory
San Diego 52, California
- Research Laboratory of Electronics
Mass. Institute of Technology
Rm. 20B-221
Cambridge 39, Mass.
- Dr. H.G. Booker
School of Electrical Engineering
Cornell University
Ithaca, N.Y.
- Department of Mathematics
Carnegie Institute of Technology
Pittsburgh, Penna.
- Signal Corps Engineering Labs.
Technical Documents Center
Evans Signal Laboratory
Falmur, N.J.
- Commanding General
Signal Corps Engineering Labs.
ATTN: Technical Reports Library
Fort Monmouth, N.J.
- Chief of Staff
HQ United States Air Force
Pentagon, Washington 25, D.C.
ATTN: AFMED - 5
- Office of Chief Signal Officer
Signal Plans and Operations Div.
Com. Liaison Br., Radio Prop. Section
The Pentagon, Washington 25, D.C.
ATTN: SIGOL - 2, Rm. 2C
- Library
Boulder Laboratory
National Bureau of Standards
Boulder, Colorado
- Naval Research Laboratory
Washington 25, D.C.
ATTN: Technical Data Section
- Air Force Development Field Rep.
Naval Research Laboratory
Code 1110
Washington 25, D.C.
- Ballistics Research Laboratory
Aberdeen Proving Ground
Aberdeen, Maryland
ATTN: Dr. Keats Pullen
- Mass. Institute of Technology
Lincoln Laboratory
P.O. Box 390
Cambridge 39, Mass.
ATTN: Dr. T.J. Carroll
- Mr. Keeve Siegel
Willow Run Research Center
Willow Run Airport
Ypsilanti, Michigan
- Willow Run Research Center
University of Michigan
Willow Run Airport
ATTN: Dr. C.L. Dolph
- Director
Naval Research Laboratory
Code 3h80
Washington 25, D.C.
ATTN: Dr. L.C. van Atta
- Director
Naval Research Laboratory
Washington 25, D.C.
ATTN: Mr. Robert E. Roterson
- National Bureau of Standards
Computation Laboratories
Washington 25, D.C.
ATTN: Dr. John Todd
- Dr. A.C. McNish
National Bureau of Standards
Washington 25, D.C.
- Dr. Terrick H. Leimer
Department of Mathematics
University of California
Berkeley, California
- Professor Bernard Epstein
Department of Mathematics
Stanford University
Stanford, California
- Division of Electrical Engineering
Electronics Research Laboratory
University of California
Berkeley 4, California
ATTN: Dr. Samuel Silver
- Dr. Joseph Kaplan
Department of Physics
University of California
Los Angeles, California
- Dr. A. Erdelyi
California Inst. of Technology
1201 E. California St.
Pasadena, California
- Dr. Robert Kalaba
Electronics Division
Rand Corporation
Santa Monica, California
- Stanford Research Institute
Stanford, California
ATTN: Dr. J.V.M. Granger
Head, Radio Systems Laboratory
- Dr. Vic Twersky
Electronics Defense Laboratory
Box 205
Mountain View, California
- Hughes Aircraft Company
Research and Development Library
Culver City, California
ATTN: M. Fodner
- Georgia Institute of Technology
Engineering Experimental Station
Atlanta, Georgia
ATTN: Dr. James E. Boyd
- Applied Physics Laboratory
The Johns Hopkins University
3642 Georgia Ave.
Silver Spring, Maryland
ATTN: Mr. F.T. McClure
- Dr. Donald E. Kerr
Department of Physics
The Johns Hopkins University
Baltimore 18, Maryland
- Dr. A. Weinstein
Institute of Fluid Dynamics and
Applied Mathematics
University of Maryland
College Park, Maryland
- Professor J.A. Pierce
Harvard University
Cambridge 38, Mass.
- Dr. F.M. Wiener
Bolt Beranek and Newman Inc.
16 Elliot St.
Cambridge 38, Mass.
- Mrs. Marjorie L. Cox, Librarian
Technical Reports Collection
Room 303A, Pierce Hall
Harvard University
Cambridge 38, Mass.
- Dr. Arthur A. Oliner
Microwave Research Institute
Polytechnic Institute of Brooklyn
55 Johnson St.
Brooklyn, N.Y.
- Professor J.H. Mulligan
School of Engineering
New York University
New York, N.Y.
- Professor Albert Heins
Carnegie Institute of Technology
Pittsburgh, Penna.

Professor Fred A. Ficken
University of Tennessee
Knoxville, Tennessee

Miss Barbara C. Grimes, Librarian
Federal Communications Commission
Washington 25, D.C.

Carnegie Institute of Washington
Department of Terrestrial Magnetism
5241 Broad Branch Road, N.W.
Washington 15, D.C.
ATTN: Library

Mathematical Reviews
190 Rye Street
Providence, Rhode Island

Mr. Martin Katzin
154 Fleetwood Terrace
Silver Spring, Maryland

Prof. R.H. Bissinger
Lebanon Valley College
Annville, Pennsylvania

Applied Physics Laboratory
The Johns Hopkins University
8621 Georgia Avenue
Silver Spring, Maryland
ATTN: Dr. B.S. Goussary

Dr. Charles H. Papas
Dept. of Electrical Engineering
California Institute of Technology
Pasadena, California

Dr. Rodman Doll
311 W. Cross St.
Ypsilanti, Michigan

California Institute of Technology
Pasadena 4, California
ATTN: Mr. Calvin Wilcox

Technical Director
Combat Development Dept.
Army Electronic Proving Ground
Fort Huachuca, Arizona

National Bureau of Standards
Boulder, Colorado
ATTN: Dr. W.R. Gallet

Mrs. A.M. Gray
Engineering Library, Plant 5
Grumman Aircraft Corp.
Bethpage, L.I., N.Y.

Professor Bernard Friedman
Dept. of Mathematics
University of California
Berkeley, California

Dr. Jane Scanlon
224-03 67th Avenue
Bayside 64, N.Y.

Dr. Solomon L. Schwebel
3689 Louis Road
Palo Alto, California

Engineering Library
University of California
405 Hilgard Avenue
Los Angeles 24, California

University of Minnesota
The University Library
Minneapolis 14, Minnesota
ATTN: Exchange Division

Lincoln Laboratory
Massachusetts Institute of Technology
P.O. Box 73
Lexington 73, Massachusetts
ATTN: Dr. Shou Chin Wang, Rm. C-351

Technical Research Group
17 Union Square West
New York 3, N.Y.
ATTN: Dr. L. Goldmuntz

Hoffman Laboratories, Inc.
Advanced Development Section
3761 South Hill Street
Los Angeles 7, California
ATTN: Dr. Richard B. Barrar

Institute of Fluid Dynamics and
Applied Mathematics
University of Maryland
College Park, Maryland
ATTN: Dr. Elliott Montroll

Brandeis University
Waltham, Massachusetts
ATTN: Library

General Electric Company
Microwave Laboratory
Electronics Division
Stanford Industrial Park
Palo Alto, California
ATTN: Library

Dr. John B. Smyth
Smyth Research Associates
3930 4th Avenue
San Diego 3, California

Dr. Georges G. Weill
Electrical Engineering
California Institute of Technology
Pasadena, California

Naval Research Laboratory
Washington 25, D.C.
ATTN: Dr. Henry J. Passerini
Code 5278 A

Dr. George Kear
5 Culver Court
Orinda, California

California Institute of Technology
Electrical Engineering
Pasadena, California
ATTN: Dr. Zohrab A. Kaprielian

Prof. Nathan Marcuvitz
Microwave Research Institute
Brooklyn Polytechnic
55 Johnson Street
Brooklyn, N.Y.

Dr. Lester Kraus
4935 Whitehaven Way
San Diego 10, California

Dr. Jerry Smoys
Dept. of Electrical Engineering
Brooklyn Polytechnic
85 Livingston Street
Brooklyn, N.Y.

Dr. I. Kolodner
Dept. of Mathematics
University of New Mexico
Albuquerque, New Mexico

Dr. Harry Hochstadt
Dept. of Mathematics
Polytechnic Institute of Brooklyn
Johnson and Jay Streets
Brooklyn, N.Y.

Hycan Eastern Inc.
75 Cambridge Parkway
Cambridge, Massachusetts
Attn: Seymour Stein

Mr. K.S. Kelleher, Section Head
Melpar, Inc.
3000 Arlington Boulevard
Falls Church, Virginia

Department of Engineering
Brown University
Providence, R.I.
ATTN: Dr. V.M. Papadopoulos

Wayne State University
Kresge-Hocker Science Library
5250 Second Boulevard
Detroit 2, Michigan

Dr. J. Brandstatten
Stanford Research Institute
S. Pasadena, California

ARRA
1 Bond Street
Westbury, L.I., N.Y.
ATTN: Dr. Norman Spector

Varian Associates
611 Hansen Way
Palo Alto, California
ATTN: (Mrs.) Perry Conway
Technical Librarian

Air Force Cambridge Research Center
Laurence G. Hanscom Field
Bedford, Massachusetts
ATTN: Dr. Ming S. Wong, CRRKP

Professor C. A. Wootton, Director
Eaton Electronics Research Laboratory
McGill University
Montreal, Canada

Dr. D.S. Jones
University of North Staffordshire
Keele, Staffordshire
England

Science Abstracts
Institute of Electrical Engineers
Savoy Place
London W.C. 2, England
ATTN: R.M. Crowther

Dr. Jean-Claude Simon
Centre de Recherches techniques
Compagnie generale de T.S.F.
Paris 19, France

Professor A. van Wijngaarden
Mathematisch Centrum
2^e Boerhaavestraat 49
Amsterdam-Zuid, Holland

Dr. C.J. Bouwkamp
Philips Research Laboratories
Eindhoven, Netherlands

Dr. E.T. Copson
Department of Mathematics
United College
University of St. Andrews
St. Andrews, Scotland

Université de Paris
Cabinet du Département
des Sciences Mathématiques
Institut Henri Poincaré
11 Rue Pierre Curie
Paris 5^e, France

Dr. W. Elwyn Williams
Dept. of Applied Math.
University of Liverpool
Liverpool, England

University of Cambridge
Cavendish Laboratory
Cambridge, England
ATTN: Prof. Philip Clemmow

Technische Hogeschool
Instituut voor Toegepaste Wiskunde
Jaffalaan 162
Delft, Holland
ATTN: Prof. R. Timman

Instituto de Fisica Teorica
Rue Pamplona, 145
S. Paulo, Brasil

Dr. H. Bremner
Philips Research Laboratories
Eindhoven, Netherlands

Electrotechnical Laboratory
Nagata-cho, Chiyoda-ku
Tokyo, Japan
ATTN: Library

Dr. A.T. De Hoop
Laboratorium voor Electrotechniek
Technische Hogeschool
Kanaalweg 2 B
Delft, Netherlands

[illegible]

Influence of edges and corners on potential

Influence of edges and corners on potential functions of surface layers...

MAY 20 1966

JUN 14 '68

**N. Y. U. Institute of
Mathematical Sciences**
25 Waverly Place
New York 3, N. Y.

